



# U.S. ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE

USAMRICD-TR-00-04

## Determination of ATP Levels in Sulfur Mustard-exposed Human Peripheral Blood Lymphocytes by a Chemiluminescent Assay

Marian R. Nelson  
Susan A. Kelly  
Clark L. Gross  
William J. Smith

January 2000

Approved for public release; distribution unlimited

DTIC QUALITY INSPECTED 4

20001226 065

U.S. Army Medical Research  
Institute of Chemical Defense  
Aberdeen Proving Ground, MD 21010-5400

## **DISPOSITION INSTRUCTIONS:**

Destroy this report when no longer needed. Do not return to the originator.

## **DISCLAIMERS:**

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

The use of trade names does not constitute an official endorsement or approval of the use of such commercial hardware or software. This document may not be cited for purposes of advertisement.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 2000		3. REPORT TYPE AND DATES COVERED May 1997 to November 1999
4. TITLE AND SUBTITLE Determination of ATP Levels in Sulfur Mustard-exposed Human Peripheral Blood Lymphocytes by a Chemiluminescent Assay			5. FUNDING NUMBERS 30162384ATC2 62384A	
6. AUTHOR(S) Nelson, MR, Kelly, SA, Gross, CL, Smith, WJ				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-DB 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-RC 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400			10. SPONSORING/MONITORING AGENCY REPORT NUMBER  USAMRICD-TR-00-04	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  Sulfur mustard (HD) is a powerful blistering agent whose mechanism is poorly understood. One of the effects of this chemical warfare agent is to lower NAD <sup>+</sup> levels through the activation of poly (ADP-ribose) polymerase; HD also depletes ATP, the major energy supply for biochemical reactions within the cell. ATP is a biological marker that can be measured in HD-exposed cells by using a chemiluminescent reaction based on the luciferin-luciferase reaction. This assay is conveniently measured by the use of a luminometer and exhibits a reproducible linear standard curve. Three prototypic inhibitors of the poly (ADP-ribose) polymerase have been studied to determine their effect on restoring or maintaining ATP levels in HD-exposed human peripheral blood lymphocytes (PBL). Two of the inhibitors, ICD 2250 (1 (2H)-isoquinolinone, 3,4-dihydro-5-[3-(methylamino)propoxy]-monohydrochloride) and ICD 2163 (3,4-dihydro-5-methyl-1(2H)-isoquinolinone), are very effective in maintaining ATP levels in HD-exposed (200 µM) PBL. With both drugs, a concentration of 100 µM restores ATP levels to nearly those found in unexposed cells with both drugs. The third inhibitor, ICD 967 (niacinamide), is not nearly as effective. This assay is now under development to test ATP levels in human epidermal keratinocytes, a more relevant <i>in vitro</i> model for vesication.				
14. SUBJECT TERMS human peripheral blood lymphocytes, sulfur mustard, HD, ATP, chemiluminescent, poly (ADP-ribose) polymerase inhibitors			15. NUMBER OF PAGES 14	
			16. PRICE CODE	
S17. ECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED
				20. LIMITATION OF ABSTRACT UNLIMITED

## Introduction

Sulfur mustard (HD) is a chemical warfare agent that causes debilitating and slowly healing injuries to eyes, lung, and skin. The mechanism(s) for causing these injuries is not currently understood, and biological markers for the severity of mustard exposure are being evaluated. HD alkylates cellular targets such as DNA, RNA, and protein (1) and depletes cellular NAD (2,3) as well as ATP, whose loss leads to metabolic disruption and the potential death of the cell (4). Maintaining or restoring ATP levels within exposed cells may lead to the amelioration of the HD lesion.

The Drug Assessment Division of the United States Army Medical Research Institute of Chemical Defense has the responsibility of determining which compounds or formulations could be potentially efficacious as medical countermeasures against chemical warfare agents. The *in vitro* team of the Basic Assessment Branch uses various relevant human *in vitro* models to develop assays for measuring critical biological markers of sulfur mustard exposure. These assays are tested using certain drugs that have the desired pharmacological actions in either prevention or treatment of the vesicant injury. One of the hypothesized consequences of mustard exposure is the loss in ATP levels and disruption of metabolic activity. A sensitive assay to measure ATP levels has been developed using a commercial BioOrbit ATP measurement kit and a Dynatech ML 3000 luminometer. The effect of HD on ATP levels in human peripheral blood lymphocytes (PBL) has been measured, and three prototypic drugs have been tested and evaluated in protecting/restoring ATP levels in these cells. Maintenance of critical ATP levels may be effective in lessening the consequences of HD exposure in a metabolically active cell.

## Materials and Methods

### *Reagents:*

RPMI 1640 media, EDTA, and trypan blue were obtained from Sigma Chemical Co., St. Louis, MO. Percoll was purchased from Pharmacia, Piscataway, NJ. Bio Orbit Monitoring Reagent Kits were obtained from Wallac Inc., Gaithersburg, MD. Trichloroacetic acid (TCA) was obtained from Wallac Oy, Turku, Finland. Sulfur mustard (CAS Registry #505-60-2, 96.8% pure) was acquired from the Edgewood Chemical and Biological Center, Aberdeen Proving Ground, MD.

### *Isolation of PBL:*

Blood was obtained by venipuncture from volunteers under an approved human use protocol. The blood was diluted with a phosphate buffered saline (PBS) (0.003M KPO<sub>4</sub>, 0.15 M NaCl, 0.1M EDTA, pH 7.4) solution (5). Thirty-five mL of diluted blood was then layered on 10 mL of Percoll (d= 1.080). The density gradient was then centrifuged at 425 xg, for 30 minutes at 20° C. The lymphocyte bands were aspirated and pooled into 50 mL centrifuge tubes and washed with Tyrode's free buffer (3). The lymphocytes were then centrifuged at 250 xg for 12 minutes at 20°C. Cell pellets were resuspended in Tyrode's free buffer, combined into one centrifuge tube, washed again in Tyrode's free

buffer and finally resuspended in RPMI 1640 buffer. The cells were counted using trypan blue on a hemacytometer and their concentration was adjusted to  $2 \times 10^7$  cells per mL.

***Exposure of lymphocytes to sulfur mustard:***

Lymphocytes were plated at  $1 \times 10^6$  cells per well in a 96-well plate. The desired pretreatment compounds were added to the appropriate wells at concentrations ranging from 200 to 2000  $\mu\text{M}$ . One-half of the plate represented control samples and samples containing buffer and pretreatment compound at designated concentrations. The other half of the plate represented lymphocyte samples to be exposed to 200  $\mu\text{M}$  HD and pretreatment compound at designated concentrations. A 10 mL stock (4000  $\mu\text{M}$ ) vial of HD was diluted 1:4 with RPMI 1640 to a working concentration of 1000  $\mu\text{M}$ , and 40  $\mu\text{L}$  of this solution was added to each well to give a final HD challenge dose of 200  $\mu\text{M}$ . After samples had been exposed to HD, the plate remained in the surety hood for one hour to allow complete hydrolysis of HD and venting of volatile agent. The plate was then placed in the 5%  $\text{CO}_2$  incubator for an additional three hours until ATP levels could be measured.

***Luminescent ATP assay:***

ATP standard solution (10  $\mu\text{M}$ ) is prepared by adding 10 mL of doubly distilled, deionized water to the vial of ATP supplied in the Bio Orbit ATP assay kit. After withdrawing enough ATP standard solution (1.0 mL) to perform a standard curve, the remaining reconstituted ATP standard is distributed into 1.0 mL aliquots in Eppendorf tubes and frozen at  $-20^\circ\text{C}$ .

**ATP Standard Curve Procedure**

Sample	Tris-Acetate mL (pH 7.7)	mL 10 $\mu\text{M}$ ATP	** mL 1 $\mu\text{M}$ ATP	[ATP]
1	0.100	0.100	-----	5.0 $\mu\text{M}$
2	0.150	0.100	-----	4.0 $\mu\text{M}$
3	0.375	0.125	-----	2.5 $\mu\text{M}$
4	3.600	0.400	-----	1.0 $\mu\text{M}$ **
5	0.250	-----	0.250	0.5 $\mu\text{M}$
6	0.225	-----	0.025	0.1 $\mu\text{M}$
7	0.190	-----	0.010	0.05 $\mu\text{M}$

Twenty-five  $\mu\text{L}$  of each of the above samples was transferred to a Micro-Lite 96-well plate, and each standard was run in triplicate.

The 96-well plate containing the HD-exposed PBL was removed from the  $37^\circ\text{C}$  incubator after three hours. Cells were solubilized by the addition of 10  $\mu\text{L}$  of 20% TCA to each well, and the plate was incubated for 15 minutes at room temperature. The samples (210  $\mu\text{L}$ ) were removed from the wells and placed in tubes containing 790  $\mu\text{L}$  of 0.2 M Tris-

acetate, pH 7.75. Aliquots of 25  $\mu\text{L}$  from each sample were pipetted (in quadruplicate) into a Micro Lite 96-well plate containing 0.125  $\mu\text{L}$  of 0.1M Tris buffer, and the plate was placed in a ML 3000 Luminometer. ATP Monitoring Reagent was reconstituted by adding 10 mL of doubly distilled deionized water to the reagent vial. This assay was performed at room temperature as recommended in the product literature (6). The luminometer was set up to inject 36  $\mu\text{L}$  of ATP Monitoring Reagent and 7  $\mu\text{L}$  of ATP standard (spike) into each well prior to reading samples.

#### *Data Analysis:*

Data was analyzed by One Way Analysis of Variance (ANOVA), and the level of significance was  $p < 0.001$  (\*) and  $p < 0.05$  (\*\*)

### **Results**

A typical standard curve of ATP is shown below in Figure 1. The curve was linear between 5 nM and 5  $\mu\text{M}$  ATP and is reproducible. The  $r^2$  value was very consistent at the 0.99 level using this chemiluminescent assay.




Figure 1. A standard curve of ATP concentrations between 50 nM and 5  $\mu\text{M}$  using the chemiluminescent ATP assay. Each point is the mean  $\pm$  s.d. value of quadruplicate determinations.

When various concentrations of HD were added to PBL, intracellular ATP levels decreased in a dose-dependent manner (Figure 2). This reduction appeared to be in two phases: a rapid initial ATP loss between 0 and 50  $\mu$ M HD, and a slower decrease that seemed to reach an approximate maximum of 50% loss at 250  $\mu$ M HD.




Figure 2. Profile of ATP levels in freshly isolated human PBL exposed *in vitro* to varying concentrations of HD. At 3 hours postexposure, ATP levels were measured using the chemiluminescence test kit (as described in Materials and Methods). Each point represents the mean value ( $\pm$  s.e.m) of three separate experiments.

When HD concentration was increased to 500  $\mu$ M, there was no further loss in ATP levels beyond that seen at 250  $\mu$ M. The challenge dose of 200  $\mu$ M HD was used in subsequent experiments since it appears from data shown in Figure 2 that the maximum effect occurs at this dose.

Inhibitors of poly (ADP-ribose) polymerase were investigated to determine whether the ATP levels of lymphocytes would be spared or restored by the addition of these compounds to the cells before they were exposed to HD. Three prototypic compounds were selected from known inhibitors of this enzyme, and the results from these studies appear below. The poly ADP-ribose polymerase (PARP) inhibitor ICD 2250 (Fig. 3) was added to the cells immediately prior to HD exposure, and ATP levels were determined at 3 hr postexposure.




Figure 3. Percent of ATP values obtained when human PBL were exposed *in vitro* to 200  $\mu$ M HD and assayed 3 hr postexposure with a chemiluminescence ATP test kit in the presence or absence of the poly ADP-ribose polymerase (PARP) inhibitor ICD 2250. Each point represents the mean  $\pm$  s.e.m value of six separate experiments. Level of significance was  $p < 0.001$  (\*) by ANOVA.

ATP levels were protected at ICD 2250 concentrations of 100  $\mu$ M or greater. At ICD 2250 concentrations of 10  $\mu$ M and 30  $\mu$ M, there was incomplete but important protection of ATP levels, 34% and 71% ATP protection, respectively. ICD 2250 had no effect on control ATP levels in unexposed lymphocytes.

Figure 4 illustrates the ATP levels of PBL measured in the presence or absence of 200  $\mu$ M HD when another PARP inhibitor, ICD 2163, was added. No effect on control ATP levels was observed in pretreated unexposed cells as shown below.






Figure 4. Human PBL exposed *in vitro* to 200  $\mu\text{M}$  HD were assayed 3 hr postexposure with a chemiluminescence ATP test kit in the presence or absence of the poly (ADP-ribose) polymerase (PARP) inhibitor ICD 2163. Each point represents the mean ( $\pm$  s.e.m.) value of three separate experiments. Level of significance was  $p < 0.001$  (\*) by ANOVA.

The results obtained with this PARP inhibitor were very similar to those of ICD 2250. At ICD 2163 concentrations of 10  $\mu\text{M}$  and 30  $\mu\text{M}$ , there was also incomplete protection of ATP levels, 51% and 81% ATP protection, respectively. No effect on control ATP levels with this inhibitor was observed, as in the case of ICD 2250.

However, the PARP inhibitor ICD 967 exhibited a different behavior, shown in Figure 5.




Figure 5. Typical representation of ATP response when human PBL were exposed *in vitro* to 200  $\mu\text{M}$  HD and assayed 3 hr postexposure with a chemiluminescence ATP test kit in the presence or absence of the poly(ADP-ribose)polymerase (PARP) inhibitor ICD 967. Each point represents the mean  $\pm$  s.e.m. of three separate experiments. Level of significance was  $p < 0.05$  (\*\*) by ANOVA.

Maximum ATP protection of approximately 40% occurred only at 1000  $\mu$ M ICD 967, five times the concentration of the challenge dose of HD. It is unknown at this time whether concentrations of ICD 967 would protect ATP levels of lymphocytes challenged with lower doses of HD.

## Conclusions

A chemiluminescent assay has been developed for ATP levels in HD-exposed human peripheral blood lymphocytes. This assay was linear within the concentrations of 50 nM and 5  $\mu$ M ATP, was very reproducible, and has a consistent  $r^2$  value of 0.99. The assay was easy to perform and can be learned in a short time by laboratory personnel.

ATP levels in PBL decrease in a dose-dependent manner when analyzed 3 hours after HD exposure. Two phases of ATP loss occur; a rapid initial phase between 0 and 50  $\mu$ M HD and a slower loss at higher concentrations of HD up to 500  $\mu$ M. This ATP loss appears to reach a plateau at approximately 50% and does not decline further with increasing HD concentration.

Breaks in DNA caused by alkylation activate the poly (ADP-ribose) polymerase that depletes the ATP levels in PBL. ATP loss and disruption of metabolic activity appears to be a very sensitive step in the injury and potential death of a cell following mustard exposure. Since ATP is involved in a myriad of metabolic activities, many critical biochemical pathways could be affected by this loss. Restoration of ATP levels and the resulting metabolic improvement may be helpful in reversing the cellular damage caused by this agent followed by recovery of the damaged cell.

With this goal in mind, we have investigated drugs that inhibit poly (ADP-ribose) polymerase. Three of the prototypic drugs that we investigated had varying effects on the ATP levels. ICD 967 showed a favorable effect only at very high levels.

However, ICD 2250 and ICD 2163 show beneficial effects in concentrations as low as 10  $\mu$ M and approach ATP levels of unexposed cells when 100  $\mu$ M concentration of drugs were used. Increasing the drug level to 300 or 1000  $\mu$ M did not appear to have any extra advantage. The results of this study give some hope to finding a medical countermeasure for the vesicant agent HD. The *in vitro* findings in PBL will be further investigated in human epidermal keratinocytes, another *in vitro* model that appears to be a more relevant target of mustard exposure. If these ATP results can be confirmed in this model, the drugs will be passed on for further testing in an *in vivo* model currently available at this institute.

This assay was very accurate but had some technical aspects that must be considered. Temperature control was critical since the temperature optimum of the luciferin-luciferase assay is 25°C. Early results were dependent on the position of the assay wells in the plate, since the assay was slowly denatured before completion of plate reading when the heating platform was at 37°C. The assay is now performed at room temperature using temperature-equilibrated reagents. We also noticed that splashing of the reagents

was occurring inside the luminometer and may have been due to using reagents that were cold and not thoroughly dissolved. This problem was solved by pre-priming injector probes with room temperature reagents (25°C) and siliconizing the probe tips.

Another problem encountered was artifactual data results from a high background thought to be due to exogenous light activation. Wrapping the ATP Monitoring Reagent syringe in aluminum foil to preclude light activation solved the problem.

## References

- (1) Papirmeister B, Feister AJ, Robinson SI, and Ford RD (1991) *Medical Defense Against Mustard Gas: Toxic Mechanisms and Pharmacological Implications*, CRC Press, Boca Raton, Florida.
- (2) Gross CL, Meier HL, Papirmeister B, Brinkley FB, and Johnson JB (1985) Sulfur Mustard Lowers Nicotinamide Adenine Dinucleotide Concentrations in Human Skin Grafted to Athymic Nude Mice, *Toxicol. Appl. Pharmacol.*, **81**, 85-90.
- (3) Meier HL, Gross CL, and Papirmeister B (1987) 2,2'-Dichlorodiethyl Sulfide (sulfur mustard) decreases NAD<sup>+</sup> levels in human leukocytes, *Toxicol. Lett.* **39**, 109-122.
- (4) Papirmeister B, Gross CL, Meier HL, Petrali JP, and Johnson JB (1985) Molecular Basis for Mustard-induced Vesication, *Fund. Appl. Toxicol.* **5**, S134-S149.
- (5) Corun CM, Kelly SA, Gross CL, and Smith WJ (1999) Development of an *In Vitro* Assay of Mustard-induced Proteolysis Using Human Cells in Culture, USAMRICD-TR-00-02. U. S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, MD, January 2000.
- (6) BioOrbit 1243-102 ATP Monitoring Kit Technical Bulletin, BioOrbit Oy, FIN-20521 Turku, Finland.

## DISTRIBUTION LIST

Addresses	Copies		
DEFENSE TECHNICAL INFORMATION CENTER ATTN DTIC OCP 8725 JOHN J KINGMAN RD STE 0944 FT BELVOIR VA 22060-6218	2	DIRECTOR ARMED FORCES MEDICAL INTELLIGENCE CENTER 1607 PORTER STREET FORT DETRICK MD 21702-5004	1
COMMANDER US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND ATTN MCMR PLD 504 SCOTT ST FORT DETRICK MD 21702-5012	2	COMMANDER US ARMY INSTITUTE OF DENTAL RESEARCH BUILDING 40 WASHINGTON DC 20307-5300	1
HQDA OFFICE OF THE SURGEON GENERAL 5109 LEESBURG PIKE SUITE 691 FALLS CHURCH VA 22041-3258	1	COMMANDER US ARMY INSTITUTE OF SURGICAL RESEARCH BUILDING 2653 FORT SAM HOUSTON TX 78234-6200	1
DIRECTOR WALTER REED ARMY INSTITUTE OF RESEARCH ATTN MCMR UWZ L 503 ROBERT GRANT AVENUE SILVER SPRING MD 20910-7500	1	COMMANDER USAMEDD CENTER & SCHOOL ATTN MCCS FC FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY AEROMEDICAL RESEARCH LABORATORY ATTN SCIENTIFIC INFORMATION CENTER PO BOX 577 FORT RUCKER AL 36362-5000	1	COMMANDER USAMEDD CENTER & SCHOOL ATTN MCCS FCD FORT SAM HOUSTON TX 78234-6100	1
COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF INFECTIOUS DISEASES 1425 PORTER ST FORT DETRICK MD 21702-5011	1	DIRECTOR ENVIRONMENTAL AND LIFE SCIENCES OFFICE OF THE DEPUTY DIRECTOR FOR RESEARCH AND ENGINEERING ROOM 3D129 WASHINGTON DC 20301-2300	1
COMMANDER US ARMY RESEARCH INSTITUTE OF ENVIRONMENTAL MEDICINE ATTN MCMR UE ZS (MS SAFRAN) BUILDING 42 NATICK MA 01760-5007	1	COMMANDER US ARMY TRAINING AND DOCTRINE COMMAND ATTN ATMD FORT MONROE VA 23651	1
COMMANDANT US ARMY CHEMICAL SCHOOL ATTN ATZN CM C FORT MCCLELLAN AL 36205	1	COMMANDER US ARMY NUCLEAR AND CHEMICAL AGENCY 7500 BACKLICK ROAD BUILDING 2073 SPRINGFIELD VA 22150-3198	1

COMMANDER US ARMY MEDICAL MATERIEL DEVELOPMENT ACTIVITY ATTN MCMR UMD 622 NEIMAN ST FORT DETRICK MD 21702-5009	1	AFOSR/NL BUILDING RM A217 BOLLING AFB DC 20332	1
EXECUTIVE OFFICER NAVAL MEDICAL RESEARCH INSTITUTE NAVAL MEDICINE COMMAND NATIONAL CAPITAL REGION BETHESDA MD 20814	1	COMMANDER US ARMY SOLDIER AND BIOLOGICAL CHEMICAL COMMAND ATTN AMSCB CI ABERDEEN PROVING GROUND MD 21010-5423	1
DEPARTMENT OF THE NAVY NAVAL POSTGRADUATE SCHOOL DUDLEY KNOX LIBRARY 411 DYER ROAD ROOM 110 MONTEREY CA 93943-5101	1	LTC RICHARD R. STOTTS BATTELLE MEMORIAL INSTITUTE JM 3 505 KING AVENUE COLUMBUS OH 43201-2695	1
USAF ARMSTRONG LABORATORY/CFTO SUSTAINED OPERATIONS BRANCH BROOKS AFB TX 78235-5000	1	COMMANDER US ARMY MEDICAL RESEARCH INSTITUTE OF CHEMICAL DEFENSE 3100 RICKETTS POINT ROAD ATTN MCMR UV ZA MCMR UV ZB MCMR UV ZS MCMR UV RC (5 copies) MCMR UV R (11 copies) MCMR UV AI W MCMR UV D MCMR UV P MCMR UV C ABERDEEN PROVING GROUND MD 21010-5425	23
DEPARTMENT OF HEALTH AND HUMAN SERVICES NATIONAL INSTITUTES OF HEALTH THE NATIONAL LIBRARY OF MEDICINE SERIAL RECORDS SECTION 8600 ROCKVILLE PIKE BETHESDA MD 20894	1		
STEMSON LIBRARY ACADEMY OF HEALTH SCIENCES BUILDING 2840 RM 106 FORT SAM HOUSTON TX 78234-6100	1		
US ARMY RESEARCH OFFICE ATTN CHEMICAL AND BIOLOGICAL SCIENCES DIVISION PO BOX 12211 RESEARCH TRIANGLE PARK NC 27709-2211	1		